

PERTURBATIONS GROWTH AND BIAS DURING ACCELERATION ^a

LUCA AMENDOLA AND DOMENICO TOCCHINI-VALENTINI

*Osservatorio Astronomico di Roma,
Viale Frascati 33, 00040 Monte Porzio Catone (Roma), Italy*

In most models of dark energy the structure formation stops when the accelerated expansion begins. In contrast, we show that the coupling of dark energy to dark matter may induce the growth of perturbations even in the accelerated regime. In particular, we show that this occurs in the models proposed to solve the cosmic coincidence problem, in which the ratio of dark energy to dark matter is constant. Moreover, if the dark energy couples only to dark matter and not to baryons, as requested by the constraints imposed by local gravity measurements, the baryon fluctuations develop a constant, scale-independent, large-scale bias which is in principle directly observable.

As it has been shown^{1,2}, an epoch of acceleration³ with a constant non-vanishing Ω_c can be realized by coupling dark matter to dark energy. In fact, a dark energy scalar field ϕ governed by an exponential potential linearly coupled to dark matter yields, in a certain region of the parameter space, an accelerated expansion with a constant ratio Ω_c/Ω_ϕ and a constant parameter of state w_ϕ , referred to as a stationary accelerated era. Similar models have been discussed elsewhere⁴. We showed⁵ that in fact the conditions of constant Ω_ϕ and w_ϕ uniquely determine the potential and the coupling of the dark energy field. The main motivation to consider a stationary dynamics is that it would solve the cosmic coincidence problem of the near equivalence at the present of the dark energy and dark matter densities, since they scale identically with time. Further theoretical motivations for coupled dark energy have been put forward^{6,1,2}.

Consider three components, a scalar field ϕ , baryons and CDM described by the energy-momentum tensors $T_{\mu\nu(\phi)}$, $T_{\mu\nu(b)}$ and $T_{\mu\nu(c)}$, respectively. General covariance requires the conservation of their sum, so that it is possible to consider a coupling such that, for instance,

$$\begin{aligned} T_{\nu(\phi);\mu}^\mu &= \sqrt{2/3}\kappa\beta T_{(c)}\phi_{;\nu} \\ T_{\nu(c);\mu}^\mu &= -\sqrt{2/3}\kappa\beta T_{(c)}\phi_{;\nu}, \end{aligned}$$

where $\kappa^2 = 8\pi G$, while the baryons are assumed uncoupled^{7,8}, $T_{\nu(b);\mu}^\mu = 0$ because local gravity constraints indicate a baryon coupling $\beta_b < 0.01$ ^{1,2,9}. Let us derive the background equations in the flat conformal FRW metric. The equations for this model have been described in another work¹⁰, in which a similar model (but with a variable coupling) was studied. We restrict ourselves to the case in which radiation has already redshifted away. For the scalar field we assume a potential $U(\phi) = U_0 e^{-\sqrt{2/3}\kappa\phi}$. The coupling β can be seen as the relative strength of the dark

^aTo appear in the proceedings of the XXXVIIth Rencontres de Moriond, "The Cosmological Model", Les Arcs, France, March 2002.

Table 1: Critical points.

Point	x	y	v	Ω_ϕ	p	w_e
a	$-\frac{\mu}{3}$	$\sqrt{1 - \frac{\mu^2}{9}}$	0	1	$\frac{3}{\mu^2}$	$\frac{2\mu^2}{9}$
b_c	$-\frac{3}{2(\mu+\beta)}$	$\frac{\sqrt{g-9}}{2 \mu+\beta }$	0	$\frac{g}{4(\beta+\mu)^2}$	$\frac{2}{3} \left(1 + \frac{\beta}{\mu}\right)$	$\frac{\mu}{\mu+\beta}$
b_b	$-\frac{3}{2\mu}$	$\frac{3}{2 \mu }$	$\sqrt{1 - \frac{9}{2\mu^2}}$	$\frac{9}{2\mu^2}$	$\frac{2}{3}$	1
c_c	$\frac{2}{3}\beta$	0	0	$\frac{4}{9}\beta^2$	$\frac{6}{4\beta^2+9}$	$1 + \frac{4\beta^2}{9}$
d	-1	0	0	1	1/3	2
e	+1	0	0	1	1/3	2
f_b	0	0	1	0	2/3	1

matter-dark energy interaction with respect to the gravitational force. The only parameters of our model are β and μ (the constant U_0 can always be rescaled away by a redefinition of ϕ). The system is best studied in the new variables^{11,8} $x = \kappa\phi'/\sqrt{6}$, $y = \frac{\kappa a}{H} \sqrt{U/3}$, and $v = \frac{\kappa a}{H} \sqrt{\rho_b/3}$, where the prime stands for derivations respect to $\log a$. The CDM energy density parameter is obviously $\Omega_c = 1 - x^2 - y^2 - v^2$ while we also have $\Omega_\phi = x^2 + y^2$, and $\Omega_b = v^2$. The system is subject to the condition $x^2 + y^2 + v^2 \leq 1$. The critical points of the resulting dynamical system are listed in Tab. 1. We denoted with $w_e = 1 + p_{tot}/\rho_{tot} = 1 + x^2 - y^2$ the total parameter of state. On all critical points the scale factor expansion is given by $a \sim \tau^{p/1-p} = t^p$, where $p = 2/(3w_e)$, while each component scales as a^{-3w_e} . In the table we also denoted $g \equiv 4\beta^2 + 4\beta\mu + 18$, and we used the subscripts b, c to denote the existence of baryons or matter, respectively, beside dark energy.

From now on we will discuss the global attractor b_c , the only critical point that may be stationary (i.e. Ω_c and Ω_ϕ finite and constant) and accelerated. On this attractor the two parameters β and μ are uniquely fixed by the observed amount of Ω_c and by the present acceleration parameter (or equivalently by $w_e = \mu/(\mu + \beta)$). For instance, $\Omega_c = 0.20$ and $w_e = 0.23$ gives $\mu = 3$ and $\beta = 10$.

Defining the perturbation variables $\delta_c = \delta\rho_c/\rho_c$, $\delta_b = \delta\rho_b/\rho_b$, $\frac{\sqrt{6}}{\kappa}\varphi = \delta\phi$, after algebraic manipulations the following second order equations for CDM and baryons are found¹² in the synchronous gauge for the wavenumber k :

$$\begin{aligned} \delta_c'' + \frac{1}{2}(4 - 3w_e - 4\beta x)\delta_c' - \frac{3}{2}\gamma\Omega_c\delta_c &= 0 \\ \delta_b'' + \frac{1}{2}(4 - 3w_e)\delta_b' - \frac{3}{2}\Omega_c\delta_c &= 0, \end{aligned}$$

where $H = d\log a/d\tau$. These equations are valid for subhorizon scales, that is $k/H \gg 1$. Important differences with respect to the equations of the standard matter-dominated case appear clearly: the friction term is modified; the constant non-vanishing dynamical term $\gamma\Omega_c$ ^{1,2,9}, which can be much larger than unity due to the extra pull of the new interaction, drives the growth of perturbation even in presence of an accelerated expansion. The quantities x, w_e and Ω_c are given in Tab.1 as functions of the fundamental parameters μ, β and $\gamma = 1 + 4\beta^2/3$. The solutions¹² are $\delta_c = a^{m_\pm}$ and $\delta_b = ba^{m_\pm}$ where

$$\begin{aligned} m_\pm &= \frac{1}{4}[-4 + 3w_e + 4\beta x \pm \Delta] \\ b_\pm &= 3\Omega_c/(3\gamma\Omega_c + 4\beta x m_\pm) \end{aligned}$$

where $\Delta^2 = (24\gamma\Omega_c + (-4 + 3w_e + 4\beta x)^2)$. The constant $b \equiv \delta_b/\delta_c \equiv b_+$ is the bias factor of the growing solution $m \equiv m_+$. The scalar field solution is $\varphi \approx (H_0 a^{(p-1)/p}/k)^2 \delta_c (\beta\Omega_c + mbx)$. For

subhorizon wavelengths φ (which is proportional to $\delta\rho_\phi/\rho_\phi$) is always much smaller than δ_c, δ_b at the present time (although it could outgrow the matter perturbations in the future if $p > 1$).

The solutions $m_\pm, b_\pm$ apply to all the critical solutions of Tab. 1 (for $\Omega_b \neq 0$ the solution can be further generalized). It is interesting to observe that for $\mu, \beta \gg 1$ the growth exponent m_+ diverges as $\mu\beta/(\mu + \beta)$: the gravitational instability becomes infinitely strong. Let us now focus on the stationary attractor b_c . Four crucial properties of the solutions will be relevant for what follows: first, the perturbations grow (i.e. $m > 0$) for all the parameters that make the stationary attractor stable; second, the baryons are antibiased (i.e. $b < 1$) for the parameters that give acceleration; third, in the $k \gg H$ limit (and in the linear regime), the bias factor is scale independent and constant in time; and fourth, the bias is independent of the initial conditions.

If one studies the perturbations in the Newtonian gauge it can be shown that a similar bias develops for the velocity fluctuations, $\theta_b = b\theta_c$, with $\theta = ik^i \delta u_i / a$, where u_i is the peculiar velocity. This could bring interesting observational consequences.

The species-dependent coupling generates a biasing between the baryon and the dark matter distributions. In contrast, the bias often discussed in literature concerns the distribution only of the very small fraction of baryons¹³ clustered in luminous bodies and it appears to depend on luminosity and type¹⁴. A first guess could be that the bulk of baryons follow the distribution of low-luminosity objects, since they contain most of the mass¹⁵. Very recently it was found¹⁶ that in the 2dFGRS catalog the average galaxy bias is close to unity, while galaxies with $L = L_*$ are slightly antibiased ($b = 0.92 \pm 0.11$) and galaxies with $L \ll L_*$ even more so. Moreover, quite remarkably, Verde et al. (2001)¹⁶ detected a scale-independent bias from 13 to 65 $h^{-1}\text{Mpc}$, scales at which our linear calculations should hold quite well. Similarly, galaxies from the IRAS-PSCz survey are also antibiased¹⁷: $b = 0.8 \pm 0.2$, a result that agrees with other estimations¹⁸. Inclusion of baryons belonging to weakly clustered objects like Lyman- α clouds can only lower the total baryon bias¹⁹. If anything, therefore, current estimates indicate $b < 1$ for the total baryon distribution. To be conservative, here we consider only very broad limits to b : since the acceleration requires antibias, we assume $0.5 < b < 1$.

In Fig. 1 we show all the various constraints. To summarize, they are: *a*) the present dark energy density $0.6 < \Omega_\phi < 0.8$; *b*) the present acceleration ($w_e < 2/3$, implying $\beta > \mu/2$); *c*) the baryon bias $1 > b > 0.5$. On the stationary attractor there is a mapping between the fundamental parameters μ, β and the observables w_e, Ω_ϕ , so one can plot the constraints on either pair of variables. It turns out that these conditions confine the parameters in the small dark shaded area, corresponding to $0.59 < w_e < 0.67$ or

$$1.1 < \beta < 1.4, \quad 2.0 < \mu < 2.6.$$

Therefore, the parameters of the stationary attractor are determined to within 20% roughly. It is actually remarkable that an allowed region exists at all. The growth rate m is approximately 0.5 in this region. For $b > 0.73$ the possibility of a stationary accelerated attractor able to solve the coincidence problem would be ruled out.

The observation of a constant, scale-independent, large-scale antibias would constitute a strong indication in favor of a dark matter-dark energy coupling and would indicate a bias mechanism well distinguishable from the hydrodynamical one. Furthermore the growth rate m is an observable quantity that can be employed to test the stationarity, for instance estimating the evolution of clustering with redshift. The combined test of b and m will be a very powerful test for the dark matter-dark energy interaction.

It is reasonable to expect that a similar baryon bias develops whenever there is a species-dependent coupling; its observation would therefore constitute a test of the equivalence principle. At the same time, the species-dependent coupling is requested to provide stationarity without conflicting with local gravity experiments. Therefore, we conjecture that the baryon bias is a strong test for all stationary dynamics.




Figure 1: Constraints on the stationary model: below the horizontal line the expansion is accelerated; in the light grey region the bias is between 0.5 and 1; between the vertical lines Ω_ϕ is within the observed range. The dark grey region is the surviving parameter space.

References

1. D. Wands , E.S. Copeland and A. Liddle, *Ann. N.Y. Acad. Sci.* **688** (1993) 647; C. Wetterich , *A&A* **301** (1995) 321.
2. L. Amendola, *Phys. Rev.* **D60** (1999) 043501 [arXiv:astro-ph/9904120].
3. A. G. Riess *et al.* [Supernova Search Team Collaboration], *Astron. J.* **116** (1998) 1009 [arXiv:astro-ph/9805201]; S. Perlmutter *et al.* [Supernova Cosmology Project Collaboration], *Astrophys. J.* **517** (1999) 565 [arXiv:astro-ph/9812133].
4. L. P. Chimento, A. S. Jakubi & D. Pavon, *Phys. Rev.* **D62** (2000) 063508; A. A. Sen and S. Sen, *Mod. Phys. Lett. A* **16** (2001) 1303 [arXiv:gr-qc/0103098]; T. Chiba, *Phys. Rev.* **D64** (2001) 103503 [arXiv:astro-ph/0106550].
5. D. Tocchini-Valentini and L. Amendola, *Phys. Rev.* **D65** (2002) 063508 [arXiv:astro-ph/0108143].
6. M. Pietroni, arXiv:hep-ph/0203085 (2002); M. Gasperini, F. Piazza and G. Veneziano, *Phys. Rev.* **D65** (2002) 023508 [arXiv:gr-qc/0108016].
7. T. Damour, G. W. Gibbons and C. Gundlach, *Phys. Rev. Lett.* **64** (1990) 123; T. Damour and C. Gundlach, *Phys. Rev.* **D43** (1991) 3873.
8. L. Amendola, *Phys. Rev.* **D62** (2000) 043511 [arXiv:astro-ph/9908023]; L. Amendola, *Phys. Rev. Lett.* **86**, 196 (2001) [arXiv:astro-ph/0006300].
9. T. Damour, arXiv:gr-qc/9606079 (1996).
10. L. Amendola and D. Tocchini-Valentini, *Phys. Rev.* **D64** (2001) 043509 [arXiv:astro-ph/0011243].
11. E. J. Copeland, A. R. Liddle and D. Wands, *Phys. Rev.* **D57** (1998) 4686 [arXiv:gr-qc/9711068].
12. L. Amendola and D. Tocchini-Valentini, arXiv:astro-ph/0111535 (2001).
13. M. Persic & P. Salucci, *MNRAS* **309** (1999) 923.
14. N. Kaiser, *Astrophys. J.* **284** (1984) L9; Bower R.G., Coles P., Frenk C.S. and White S.D.M., *Ap.J.* **405** (1993) 403; J. A. Peacock and S. J. Dodds, *Mon. Not. Roy. Astron. Soc.* **267** (1994) 1020 [arXiv:astro-ph/9311057]; Benoist C. et al., *Astrophys. J.* **472** (1996) 452.
15. J. Loveday, *Astrophys. J.* **489** (1997) 22; P. Valageas, J. Silk & R. Schaeffer, *A&A* **366** (2001) 363.
16. L. Verde et al., arXiv:astro-ph/0112161 (2001).
17. H. A. Feldman, et al., *Phys. Rev. Lett.* **86** (2001) 1434.
18. K. Rines, M.J. Geller, M.J. Kurtz, A. Diaferio, T.H. Jarrett, J.P. Huchra, arXiv:astro-ph/0109425 (2001).

19. M. Douspis, A. Blanchard, & J. Silk, arXiv:astro-ph/0110181 (2001).